MycoKeys 106: 201–224 (2024)
DOI: 10.3897/mycokeys.106.125831

Research Article

Morphological characteristics and phylogenetic analyses revealed four new wood inhabiting fungi (Agaricomycetes, Basidiomycota) in Xizang Autonomous Region, China

Hong-Min Zhou[1,2], Xun-Chi Zhang[2], Jie-Ting Li[3], Fang Wu[4], Chang-Lin Zhao[1,2]

1 *The Key Laboratory of Forest Resources Conservation and Utilization in the Southwest Mountains of China Ministry of Education, Key Laboratory of National Forestry and Grassland Administration on Biodiversity Conservation in Southwest China, Yunnan Provincial Key Laboratory for Conservation and Utilization of In-forest Resource, Southwest Forestry University, Kunming 650224, China*
2 *College of Forestry, Southwest Forestry University, Kunming 650224, China*
3 *Key Laboratory of Forest Ecology in Tibet Plateau, Ministry of Education, Institute of Tibet Plateau Ecology, Tibet Agricultural & Animal Husbandry University, Nyingchi, Tibet 860000, China*
4 *Institute of Microbiology, School of Ecology and Nature Conservation, Beijing Forestry University, Beijing 100083, China*
Corresponding authors: Chang-Lin Zhao (fungichanglinz@163.com); Fang Wu (fangwubjfu2014@yahoo.com)

Abstract

Four new fungi from Xizang in southwest China, *Calocera ramaria*, *Ceraceomyces rhizomorphus*, *Leptosporomyces linzhiensis*, and *Ramaria xizangensis* are described and illustrated based on the morphological and molecular evidence. *Calocera ramaria* is characterized by the ramal and bright orange basidiomata, a monomitic hyphal system with simple septa generative hyphae, usually 4-septate basidiospores; *Ceraceomyces rhizomorphus* is characterized by the cream to yellowish basidiomata with rhizomorphs, cylindrical basidiospores; *Leptosporomyces linzhiensis* is characterized by white with pink basidiomata, cylindrical to oblong ellipsoid basidiospores; *Ramaria xizangensis* is characterized by flesh pink basidiomata, branched dichotomously in 4–5 ranks, a monomitic hyphal system with clamped generative hyphae, ellipsoid to cylindrical and densely warted basidiospores.

Key words: Molecular systematic, phylogenetic analysis, taxonomy, wood-decaying fungi

OPEN ACCESS

This article is part of:
Diversity, taxonomy, and systematics of macrofungi from tropical Asia
Edited by Olivier Raspé, Rui-Lin Zhao, Jennifer Luangsa-ard

Academic editor: Rui-Lin Zhao
Received: 20 April 2024
Accepted: 27 May 2024
Published: 25 June 2024

Citation: Zhou H-M, Zhang X-C, Li J-T, Wu F, Zhao C-L (2024) Morphological characteristics and phylogenetic analyses revealed four new wood inhabiting fungi (Agaricomycetes, Basidiomycota) in Xizang Autonomous Region, China. MycoKeys 106: 201–224. https://doi.org/10.3897/mycokeys.106.125831

Copyright: © Hong-Min Zhou et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The fruiting bodies of Basidiomycota exhibit complex forms, such as gilled, poroid, toothed, coralloid basidiomata. Numerous taxonomists have endeavored to construct a stable classification system based on these characters (Gäumann 1953). Recently, the analysis of DNA sequences has emerged as a common method for deducing fungal phylogenies and enhancing higher classification frameworks through the integration of genetic traits (Cui et al. 2019; Wijayawardene et al. 2020; Liu et al. 2023).

The abundance of biodiversity in *Abies* forests can be attributed to the plentiful presence of humus and mycorrhizal fungi, which foster an optimal environment for the proliferation of the macrofungal species. Information regarding the fungal diversity in *Abies* communities is scattered over a range of publications

(Ryvarden and Gilbertson 1993; Dai 2022). *Ceraceomyces* Jülich, a small genus characterized by yellow rhizomorphic basidiomata, was established by Jülich based on the taxon *C. tessulatus* (Cooke) Jülich (Jülich 1972). This genus, originally from North America, features annual, resupinate, pellicular basidiomata with a smooth or merulioid hymenial surface, a monomitic hyphal system, narrowly clavate basidia, and subglobose, narrowly ovate, ellipsoid to cylindrical basidiospores (Chikowski et al. 2016). Phylogenetic studies revealed that *Ceraceomyces* was polyphyletic, comprising three distinct groups. The section of *Corticium tessulatum* Cooke belonged to Polyporales, and *Ceraceomyces serpens* (Tode) Ginns and *C. eludens* K.H. Larss. were part of phlebioid clade (Larsson et al. 2004). A recent study indicated that the type species, *Corticium tessulatum* is classified under the order Amylocorticiales (Binder et al. 2010; Chikowski et al. 2016), as well as species, *C. yunnanensis* Qi Yuan & C.L. Zhao and *C. borealis* (Romell) J. Erikss. & Ryvarden (Yuan et al. 2023). Currently, eleven species are recognized in the genus *Ceraceomyces*, including *C. cystidiatus* (J. Erikss. & Hjortstam) Hjortstam, *C. eludens*, *C. microsporus* K.H. Larss. and *C. sublaevis* (Bres.) Jülich were accepted in the genus. A genus, *Crystallicutis* El-Gharabawy, Leal-Dutra & G.W. Griff. was derived from *Ceraceomyces* based on the crystals in the hymenium and subiculum of the basidiomata, which includes the brown-rot species *C. serpens* (El-Gharabawy et al. 2021). Both species *C. sulphurinus* and *C. violascens* (Fr.) Jscens were recorded in *Ceraceomyces*, are considered congeneric with *Rhizochaete* Gresl., Nakasone & Rajchenb. due to the characteristics like the rhizomorphic margin and the purple reaction in KOH.

Calocera (Fr.) Fr. is known for its distinctive characteristics, stipitate, fasciculate or scattered, gelatinous basidiomata, dendroid or staghorn-like, subclavate to clavate basidia and probasidia, as well as cylindrical to reniform, septate or non-septate basidiospores (Fisher 1931; Lowy 1971; Peng et al. 1992; Wu et al. 2011; Shirouzu et al. 2017). Recent phylogenetic analyses of the class Dacrymycetes demonstrated that *Calocera* was polyphyletic and species in the genus are scattered throughout the family Dacrymycetaceae together with most of the species of *Dacrymyces* Nees (1817: 89) as well as a few species from other genera such as *Dacryopinax* G.W. Martin (1948: 116) and *Femsjonia* Fr. (Shirouzu et al. 2007; Zamora and Ekman 2020).

The genus *Leptosporomyces* Jülich is characterized by the resupinate basidiomata, white yellow and smooth hymenial surface, a monomitic hyphal system with clamped connections, and thin-wall, smooth, acyanophilous basidiospores. Recent research has indicated that *Leptosporomyces* was polyphyletic, with two taxa, *L. galzinii* (Bourdot) Jülich and *L. raunkiaeri* (M.P. Christ.) Jülich, grouped in the order Atheliales, while *L. septentrionalis* (J. Erikss.) Krieglst. was placed in the order Amylocorticiales (Larsson 2007; Hodkinson et al. 2014; Sulistyo et al. 2021). The generic delimitation of *Fibulomyces* Jülich and *Leptosporomyces* remains controversial, with both being indistinguishable in phylogenetic and morphological analyses, leading to the former being considered as a synonym of the latter (Bernicchia and Gorjón 2010).

Ramaria Fr. ex Bonord. is a widely distributed non-gilled Basidiomycete genus (Marr and Stuntz 1973; Petersen 1981; Humpert et al. 2001). The genus is recognized by branched basidiomata, mono- to dimitic hyphal systems with clamped or simple-septate generative hyphae, and smooth to echinulate,

verrucose-reticulate or striate ornamentation basidiospores (Corner 1950; Marr and Stuntz 1973; Petersen 1981; Humpert et al. 2001). The genus has been classified into four subgenera, namely *R.* subg. *Ramaria*, *R.* subg. *Laeticolora* Marr & D.E. Stuntz, *R.* subg. *Lentoramaria* Corner, and *R.* subg. *Echinoramaria* Corner (Marr and Stuntz 1973; Humpert et al. 2001; Exeter et al. 2006; Knudson 2012; Hanif et al. 2019). Initially, *Ramaria* was treated as a subgenus within *Clavaria* (Coker 1923; Doty 1944) until Corner (1970) elevated it to genus rank. Studies based on the morphological and molecular data agree on the paraphyletic state of *Ramaria* (Humpert et al. 2001; Hosaka et al. 2006; Giachini et al. 2010).

In the present paper, species from four genera are collected from Xizang under forest of *Abies*, and the phylogenetic relationships of four taxa are still unclear. Thus, to explore the diversity and taxonomic status with different characters for those taxa will be significant for macrofungi in Xizang, and the taxonomy and phylogeny analysis show that they are new to science.

Material and method

The specimens were collected from Xizang which were deposited in the herbarium of the Southwest Forestry University (SWFC), Kunming, Yunnan Province, China. Samples were photographed when fresh in the field, and their habitats were recorded. Microscopic structures were discussed by Zhao et al. (2023). Special color terms were set by Anonymous (1969) and Petersen (1996). A Nikon Digital Sight DS-L3 or Leica ICC50 HD camera (magnification ×1,000) was used to exam hand-cut sections of basidiomata, which were first treated with 5% KOH for a few minutes and then with 1% phloxine B ($C_{20}H_4Br_4Cl_2K_2O_5$). At least 30 basidiospores of each species were examined. The values were expressed as a mean with 5% of the measurements excluded from each end of the range, given in parentheses. Stalks were excluded for basidia measurement, and the hilar appendages were excluded for basidiospore measurement.

DNA extraction, amplification, and sequencing

The CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd., Beijing) was used to obtain DNA from dried specimens and PCR was performed according to the manufacturer's instructions with some modifications (Yang et al. 2023). ITS were amplified using the primer pairs ITS5/ITS4 (White et al. 1990). The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min, followed by 35 cycles at 94 °C for 40 s, 54 °C for 45 s, and 72 °C for 1 min; and a final extension at 72 °C for 10 min. The PCR procedure for LSU was as follows: initial denaturation at 94 °C for 1 min, followed by 35 cycles at 94 °C for 30 s, 50 °C for 1 min, and 72 °C for 1.5 min; and a final extension at 72 °C for 10 min. All newly generated sequences were submitted to GenBank and are listed in Table 1.

Sequences generated for this study were aligned, with additional sequences downloaded from GenBank. Sequences were aligned using MAFFT v.7 (https://mafft.cbrc.jp/alignment/server/), adjusting the direction of nucleotide sequences according to the first sequence (accurate enough for most cases), and selecting the G-INS-i iterative refinement method (Katoh et al. 2019). Alignments were

Table 1. Taxa information and the sequences used in this study. *Newly generated sequences for this study.

Species	Locality	Voucher	ITS	LSU
Amyloathelia crassiuscula	Sweden	GB/K169-796	DQ144610	–
Amylocorticium cebennense	USA	HHB-2808	GU187505	GU187561
Amylocorticium subincarnatum	Sweden	GB/AS-95	AY463377	AY586628
Amylocorticium subsulphureum	USA	HHB-13817	GU187506	GU187562
Anomoporia bombycina	USA	CFMR L-6240	GU187508	GU187611
Anomoporia vesiculosa	China	Dai 22795	ON413718	ON413720
Athelia abscondita	USA	Goyette 633	OP877120	OP902328
Athelopsis subinconspicua	Sweden	GB0058732	LR694197	LR694174
Bondarzewia occidentalis	Canada	AFTOL-ID 452	DQ200923	DQ234539
Byssocorticium caeruleum	Canada	RS 09400 (H)	NR_121454	–
Calocera bambusicola	China	Wu 9910-12	FJ195751	–
Calocera cornea	Sweden	UPS F 940775	MN595627	MN595627
Calocera cornea	Unknown	AFTOL ID 438	AY789083	AY701526
Calocera cornea	Sweden	UPS F 940774	MN595626	MN595626
Calocera cornea	Canada	CBS 124.84	AB712437	AB472738
Calocera guepinioides	New Zealand	PDD 107969	LC131411	LC131370
Calocera guepinioides	New Zealand	PDD 107981	LC131412	LC131371
Calocera guepinioides	New Zealand	PDD 105005	LC131407	LC131366
Calocera guepinioides	New Zealand	PDD 107874	LC131409	LC131368
Calocera guepinioides	New Zealand	PDD 105033	LC131408	LC131367
Calocera guepinioides	New Zealand	PDD 107929	LC131410	LC131369
Calocera lutea	New Zealand	PDD 107841	LC131413	LC131372
Calocera lutea	New Zealand	PDD 107842	LC131414	LC131373
Calocera palmata	New Zealand	PDD 107830	LC131415	LC131374
Calocera palmata	New Zealand	PDD 107925	LC131416	LC131375
Calocera palmata	Japan	CBS 127.51	MH856777	MH868295
Calocera ramaria	China	CLZhao 31166	**PP399147**	**PP862915**
Calocera sinensis	China	MHHNU30743	MK167408	–
Calocera sinensis	China	Wu 0505-3	FJ195753	–
Calocera sinensis	China	Wu 0703-6	FJ195754	–
Calocera sinensis	China	JCH 070726	FJ195755	–
Calocera tibetica	China	Dai 20171	MW549777	MW750403
Calocera tibetica	China	Dai 20178	MW549778	MW750404
Calocera viscosa s.lat.	Sweden	UPS F-940773	MN595628	MN595628
Calocera viscosa s.lat.	Germany	FTOL ID1679	DQ520102	DQ520102
Ceraceomyces americanus	USA	FP-102188	KP135409	KP135277
Ceraceomyces atlanticus	Brazil	URM 85888	NR_153926	NG_060427
Ceraceomyces atlanticus	China	M67	OR766067	KX685874
Ceraceomyces borealis	Sweden	KHL 8432	EU118610	–
Ceraceomyces eludens	Sweden	JS 27108	AF090879	–
Ceraceomyces eludens	Sweden	JS 22780	AF090877	–
Ceraceomyces eludens	United Kingdom	KM 194563	OR907143	–
Ceraceomyces microsporus	USA	UC 2023077	KP814418	–
Ceraceomyces microsporus	Sweden	JS 27153	AF090873	–
Ceraceomyces rhizomorphus	China	CLZhao 31154	**PP399151**	–
Ceraceomyces rhizomorphus	China	CLZhao 31161	**PP399148**	–
Ceraceomyces rhizomorphus	China	CLZhao 31188	**PP399149**	**PP862917**
Ceraceomyces rhizomorphus	China	CLZhao 31197	**PP399150**	**PP862916**
Ceraceomyces sublaevis	USA	FP-101245-Sp	KP135029	GU187607
Ceraceomyces tessulatus	USA	MPN 152885038	OR680647	–
Ceraceomyces tessulatus	Sweden	KHL 16429	KU518951	–
Ceraceomyces yunnanensis	China	CLZhao 18992	OQ132519	OQ147003
Clavariadelphus amplus	China	HMAS 250466	MK705858	MK704448
Coniophora marmorata	Belgium	MUCL: 31667	GU187515	GU187571
Dacrymyces longistipitatus	New Zealand	PDD 107996	LC131425	LC131386
Dacrymyces pachysporus	New Zealand	PDD 105004	LC131429	LC131392

Species	Locality	Voucher	ITS	LSU
Dacrymyces parastenosporus	New Zealand	PDD104960	LC131431	LC131394
Dacrymyces stillatus	Sweden	UPS F-939814	MN595676	MN595676
Dacrymyces subalpinus	Japan	TUFC12834	AB712465	AB299060
Dacryonaema macnabbii	Sweden	UPS F-940949	MN595650	MN595650
Dacryonaema macnabbii	Sweden	UPS F-940951	MN595651	MN595651
Dacryonaema macrosporum	Norway	O 160045	MN595659	MN595659
Dacryonaema macrosporum	Finland	UPS F-940998	MN595660	MN595660
Dacryonaema rufum	Sweden	UPS F-941003	MN595645	MN595645
Dacryonaema rufum	Sweden	UPS F-941005	MN595646	MN595646
Dendrdacrys brasiliense	Brazil	INPA:241458	AB744230	AB723514
Dendrdacrys dendrocalami	Japan	TUFC 13914	AB712453	AB712428
Fibulomyces mutabilis	Germany	HG-B 5753 (GB)	GQ162817	–
Ganoderma resinaceum	Unknown	C45	KX371982	KX372027
Gautieria parksiana	USA	SNF 236 USA	AF377059	–
Gloeocantharellus neoechinosporus	China	GDGM 75321	MK358820	MK358815
Go. ludovicianus	USA	TFB 14476	KJ655570	KJ655580
Gomphus clavatus	Spain	MA-Fungi 48085	AJ292292	–
Hypochniciellum subillaqueatum	Sweden	KHL 8493	AY463431	AY586679
Hypochniciellum subillaqueatum	UK	KM165142	MZ159402	–
Kavinia himantia	USA	CFMR: DLL2011-079	KJ140598	–
Kavinia alboviridis	USA	CFMR: DLL2011-131	KJ140634	–
Lentaria micheneri	USA	RRD6 (TENN)	MF773634	–
Lactarius sp.	New Zealand	PDD:113066	MW683864	MW683691
Lentaria byssiseda	USA	TENN 61159	FJ596788	–
Leptosporomyces fuscostratus	USA	UC 2022884	KP814350	–
Leptosporomyces fuscostratus	Unknown	DK 16_251	OL436970	–
Leptosporomyces galzinii	Sweden	GB 0107211	LR694202	LR694180
Leptosporomyces galzinii	USA	UC 2023126	KP814291	–
Leptosporomyces linzhiensis	China	CLZhao 31174	**PP399152**	**PP862922**
Leptosporomyces linzhiensis	China	CLZhao 31183	**PP399153**	**PP862918**
Leptosporomyces linzhiensis	China	CLZhao 31187	**PP399154**	–
Leptosporomyces linzhiensis	China	CLZhao 31190	**PP399155**	–
Leptosporomyces raunkiaeri	USA	UC 2023053	KP814293	–
Leptosporomyces raunkiaeri	USA	CFMR: HHB-7628	GU187528	GU187588
Leptosporomyces septentrionalis	USA	UC 2023047	KP814348	–
Leptosporomyces septentrionalis	Sweden	GB 0090937	LR694203	LR694181
Leptosporomyces septentrionalis	Norway	JS 16122	GU187497	–
Lobulicium occultum	Sweden	KHL13496b	MT340827	–
Mythicomyces corneipes	Unknown	AFTOL-972	DQ404393	AY745707
Phaeoclavulina flaccida	Italy	AMB n. 17671	MK796107	MK796156
Phlebiella christiansenii	Finland	KHL 11689	EU118659	–
Phlebiella vaga	Sweden	KHL 11065	EU118660	EU118661
Piloderma fallax	Finland	CFMR: S-12	GU187535	–
Plicaturopsis crispa	China	LWZ 20201017-11	ON897938	ON885398
Plicaturopsis crispa	Brazil	URM 85888	NR_153926	NG_060427
Ramaria abietina	USA	u066	KY510818	–
Ramaria acrisiccescens	USA	OSC 112057	KY354738	KY354711
Ramaria admiratia	USA	TENN: 69114	NR_137862	NG_059504
Ramaria amyloidea	USA	OSC 69891	EU837196	KP637036
Ramaria apiculata var. *brunnea*	USA	CBS:149.74	MH860840	MH872577
Ramaria araiospora	Germany	OSC 108707	EU846298	–
Ramaria aurantiisiccescens	USA	OSC 104868	EU837197	–
Ramaria aurea	Italy	AMB 18352	MN637783	MN637796
Ramaria botrytis	Italy	AMB n. 18201	NR_189799	NG_241889
Ramaria botrytis	Argentina	GM 19044	OP177707	OP177871
Ramaria botrytis	USA	snf213	AF377055	–
Ramaria botrytis f. *musicolor*	Italy	ZT Myc 57160	KY626144	–
Ramaria botrytis var. *aurantiiramosa*	USA	OSC 140667	JX310410	–

Species	Locality	Voucher	ITS	LSU
Ramaria botrytis var. *aurantiiramosa*	USA	WTU-F-043053	KX574471	–
Ramaria celerivirescens	USA	OSC 140471	JX310392	JX269125
Ramaria claviramulata	USA	WTU-F-043055	KX574472	KX671009
Ramaria conjunctipes	USA	OSC: 110613	KC346861	–
Ramaria coulterae	USA	OSC 69929	EU669320	EU669320
Ramaria dendrophora	Argentina	GM 20020	OP177716	OP177880
Ramaria dendrophora	Argentina	GM 19094	OP177715	OP177879
Ramaria fennica	Italy	AMB n. 17522	MK682678	–
Ramaria flavescens	Italy	AMB 17404	KY354743	–
Ramaria flavescens	Italy	AMB 17404	MK493036	–
Ramaria flava	Italy	AMB 17393	MK493035	–
Ramaria flavinedulis	Argentina	GM 19056	OP177717	OP177881
Ramaria flavinedulis	Argentina	GM 19035	OP177720	OP177884
Ramaria flavobrunnescens var. *aromatica*	USA	AGK 059	JQ408240	–
Ramaria foetida	USA	AGK 058	JQ408239	JQ408239
Ramaria formosa	USA	OSC1064203	EU525994	–
Ramaria fumosiavellanea	USA	WTU-F-063048	MK169345	–
Ramaria gelatiniaurantia	USA	OSC 65737	KP658144	–
Ramaria inedulis	Chile	12648	OP177723	OP177887
Ramaria inedulis	Argentina	GM 19047	OP177722	OP177886
Ramaria largentii	USA	OSC 67012	KP658130	KP637058
Ramaria luteovernalis	Italy	MCVE 28637	NR_155720	KT357477
Ramaria maculatipes	USA	OSC 112051	KY354749	KY354721
Ramaria magnipes	USA	WTU-F-063057	MK169351	MK493050
Ramaria myceliosa	USA	AGK 035	JQ408230	–
Ramaria obtusissima	USA	TFB 14473	KJ655554	KJ655575
Ramaria patagonica	Argentina	403	OP177710	OP177874
Ramaria patagonica	Argentina	GM 19106	OP177713	OP177877
Ramaria pseudoflava	Italy	AMB 17392	MK493046	–
Ramaria rasilisporoides	Pakistan	MH-2013	MG760613	–
Ramaria rasilisporoides	USA	WTU-F-043029	MK169346	–
Ramaria rubella	USA	OSC 115946	EU669317	EU669343
Ramaria rubella f. *rubella*	USA	AGK 049	JQ408236	–
Ramaria rubribrunnescens	USA	OSC 119676	EU652352	EU652387
Ramaria rubribrunnescens	USA	OSC 66051	KY354750	KY354722
Ramaria sandaracina var. *sandaracina*	Canada	UBC F28386	KP454028	–
Ramaria sp.	India	KD-14-006	KT824242	–
Ramaria stricta	Germany	CBS 165.48	MH856299	–
Ramaria stricta var. *concolor*	USA	AGK 011	JQ408221	–
Ramaria stuntzii	USA	OSC 73315	KP658122	KP637048
Ramaria subbotrytis	Spain	MA-Fungi 48088	AJ408361	–
Ramaria subtilis	Spain	MA-Fungi 48055	AF442098	–
Ramaria suecica	USA	OSC 115933	KP658148	KP637079
Ramaria testaceoflava	USA	OSC 107885	KP658128	AY586708
Ramaria verlotensis	USA	WTU-F-063047	KX574480	KX671016
Ramaria xizangensis	China	CLZhao 31169	**PP399156**	**PP862919**
Ramaria xizangensis	China	CLZhao 31180	**PP399157**	**PP862920**
Ramaria xizangensis	China	CLZhao 31204	**PP399158**	**PP862921**
Ramaria formosa	Italy	AMB 18529	MT055910	MT053203
Ramaricium polyporoideum	USA	TENN: 065654	MF992160	MF992160
Stereopsis vitellina	Sweden	F 703241	LR694211	LR694189
Turbinellus floccosus	USA	MO 285170	MN319564	MN319563
Unilacryma unispora	Sweden	UPS F 941268	MN595672	MN595672
Unilacryma unispora	Sweden	UPS F 941277	MN595665	MN593500
Xenasmatella ardosiaca	Costa Rica	KHL 12928	EU118658	–
Xenasmatella ardosiaca	USA	CBS 126045	MH864060	MH875515

manually adjusted to maximize alignment and minimize gaps with BioEdit v.7.0.9 (Hall 1999). A dataset of concatenated ITS and LSU sequences was used to determine the phylogenetic position of the new species. Maximum likelihood (ML) analysis was performed using the CIPRES Science Gateway (Miller et al. 2010) based on the dataset using the RA × ML-HPC BlackBox tool, with setting RA × ML halt bootstrapping automatically and 0.25 for maximum hours and obtaining the best tree using ML search. Other parameters in ML analysis used default settings, and statistical support values were obtained using nonparametric bootstrapping with 1,000 replicates. Bayesian inference (BI) analysis based on the dataset was performed using MrBayes v.3.2.6 (Ronquist and Huelsenbeck 2012). The best substitution model for the dataset was selected by ModelFinder (Kalyaanamoorthy et al. 2017) using a Bayesian information criterion, and the model was used for Bayesian analysis. Four Markov chains were run from random starting trees. Trees were sampled every 1,000th generation. The first 25% of sampled trees were discarded as burn-in, whereas other trees were used to construct a 50% majority consensus tree and for calculating Bayesian posterior probabilities (BPPs). The aligned sequences were deposited in TreeBase (https://www.treebase.org/treebase-web/home.html; submission ID 31437).

Branches of the consensus tree that received bootstrap support for ML were greater than or equal to 75%, Bayesian posterior probabilities more than 0.9, respectively.

Result

The Phylogeny of *Calocera*

BI analysis yielded a similar topology to MP and ML analysis. Only the MP tree is provided here (Fig. 1). Branches that received bootstrap support for ML (ML-BS), and BI (BPP) greater than or equal to 75% (MP-BS and ML-BS) and 0.90 (BPP) were considered as significantly supported, respectively. The ITS and LSU dataset contains sequences from 26 fungal specimens representing twelve *Calocera* taxa. The average SD of split frequencies in BI analyses is 0.005504 (BI). The phylogenetic tree (Fig. 1) reveals the new species has close relationship with *C. tibatica*, sister to *C. viscosa* and *C. cornea*.

The Phylogeny of *Ceraceomyces*

The dataset included ITS and LSU from 29 samples representing 22 taxa. The best model for the concatenated ITS+LSU dataset estimated and applied for BI analysis was “GTR+I+G4”, datatype = DNA, nucmodel = 4by4, lset nst = 6, rates = invgamma; state frequencies had a Dirichlet prior (1,1,1,1), and the distribution was approximated using four categories. BI analysis yielded a similar topology to ML analysis, with an average standard deviation of split frequencies of 0.006593. The ML tree was provided (Fig. 2). Branches that received bootstrap support for ML and BI ≥ 70%, and 0.75 were considered significantly supported, respectively.

The analysis reveals four clades (Fig. 2), in which three European species *C. eludens*, *C. microsporus*, *C. sublaevis* clustered together and *Rhizochaete americanus* (Nakasone, C.R. Bergman & Burds.) Gresl., Nakasone & Rajchenb. The core clade formed by *C. tessulatus* and *C. atlanticus*, along with



Figure 1. Phylogeny of species in *Calocera* generated by maximum likelihood based on ITS+LSU sequence data. Branches are labeled with maximum likelihood bootstrap ≥ 75% and Bayesian posterior probabilities ≥ 0.90, respectively. New species are in bold.

Hypochniciellum subillaqueatum (Litsch.) Hjortstam. Four specimens from China formed two lineages, namely *Ceraceomyces rhizomorphus* with *C. yunnanensis*, and were sister to *C. borealis*.

The Phylogeny of *Leptosporomyces*

BI analysis yielded a similar topology to MP and ML analysis, with an average standard deviation of split frequencies = 0.008841. Only the MP tree is provided here (Fig. 3). Branches that received bootstrap support for ML (ML-BS), and BI (BPP) greater than or equal to 75% (MP-BS and ML-BS) and 0.90 (BPP) were considered as significantly supported, respectively. Four previously accepted species, *L. galzinii*, *L. fuscostratus* (Jülich) Krieglst., *L. raunkiaeri*, and *L. mundus* (H.S. Jacks. & Dearden) Jülich received strong support in three lineages. The new species *L. linzhiensis* had a close relationship with *L. septentrionalis* with full support.


OR907143 Ceraceomyces eludens KM 194563
AF090877 Ceraceomyces eludens JS 22780
AF090879 Ceraceomyces eludens JS 27108
AF090873 Ceraceomyces microsporus JS 27153
KP814418 Ceraceomyces microsporus UC 2023077
KP135029 Ceraceomyces sublaevis FP 101245 Sp
KP135409 Rhizochaete americanus FP 102188
EU118658 Xenasmatella ardosiaca KHL 12928
MH864060 Xenasmatella ardosiaca CBS 126045
EU118659 Phlebiella christiansenii KHL 11689
OQ371469 Xenasmatella bambusicola CLZhao 10985
EU118660 Phlebiella vaga KHL 11065
PP399150 Ceraceomyces rhizomorphus CLZhao 31197
PP399148 Ceraceomyces rhizomorphus CLZhao 31161
PP399149 Ceraceomyces rhizomorphus CLZhao 31188
PP399151 Ceraceomyces rhizomorphus CLZhao 31154
EU118610 Ceraceomyces borealis KHL 8432
OQ132519 Ceraceomyces yunnanensis CLZhao 18992
DQ144610 Amyloathelia crassiuscula GB/K 169-796
OR766067 Plicaturopsis crispa M 67
MZ159402 Hypochniciellum subillaqueatum KM 165142
AY463431 Hypochniciellum subillaqueatum KHL 8493
ON413718 Anomoporia vesiculosa Dai 22795
KU518951 Ceraceomyces tessulatus KHL 16429
OR680647 Ceraceomyces tessulatus MPN 152885038
NR 153926 Ceraceomyces atlanticus URM 85888
KX685875 Ceraceomyces atlanticus URM 85888
GU187506 Amylocorticium subsulphureum HHB 13817
GU187505 Amylocorticium cebennense HHB 2808
0.05

Figure 2. Phylogeny of species in *Ceraceomyces* generated by maximum likelihood based on ITS+LSU sequence data. Branches are labeled with maximum likelihood bootstrap ≥ 75% and Bayesian posterior probabilities ≥ 0.90, respectively. New species are in bold.


KP814291 Leptosporomyces galzinii UC2023126
LR694202 Leptosporomyces galzinii GB0107211
GQ162814 Byssocorticium caeruleum RS 09400
KP814293 Leptosporomyces raunkiaeri UC2023053
GU187528 Leptosporomyces raunkiaeri CFMR HHB 7628
LR694197 Athelopsis subinconspicua GB0058732
GU187535 Piloderma fallax CFMR:S-12
LR694211 Stereopsis vitellina F703241
KP814350 Leptosporomyces fuscostratus UC2022884
OL436970 Leptosporomyces fuscostratus DK16 251
GQ162817 Fibulomyces mutabilis HG-B 5753 (GB)
OP877120 Athelia abscondita SG524
MT340827 Lobulicium occultum KHL13496b
GU187515 Coniophora marmorata MUCL 31667
DQ404393 Mythicomyces corneipes AFTOL-ID 972
LT716026 Lepiota cristata ZRL 20151133
GU187497 "Leptosporomyces septentrionalis" JS16122
PP399152 Leptosporomyces linzhiensis CLZhao 31174
PP399155 Leptosporomyces linzhiensis CLZhao 31190
PP399153 Leptosporomyces linzhiensis CLZhao 31183
PP399154 Leptosporomyces linzhiensis CLZhao 31187
KP814348 Leptosporomyces septentrionalis UC2023047
LR694203 Leptosporomyces septentrionalis GB0090937
GU187505 Amylocorticium cebennense CFMR HHB:2808
GU187506 Amylocorticium subsulphureum CFMR HHB:13817
DQ200923 Bondarzewia occidentalis AFTOL-ID 452
0.04

Figure 3. Phylogeny of species in *Leptosporomyces* generated by maximum likelihood based on ITS+LSU sequence data. Branches are labeled with maximum likelihood bootstrap ≥ 75% and Bayesian posterior probabilities ≥ 0.90, respectively. New species are in bold.

The Phylogeny of *Ramaria*

BI analysis yielded a similar topology to MP and ML analysis. Only the MP tree is provided here (Fig. 4). Branches that received bootstrap support for ML (ML-BS), and BI (BPP) greater than or equal to 75% (MP-BS and ML-BS) and 0.90 (BPP) were considered as significantly supported, respectively. Four clades were obtained from our phylogenetic analysis, *Ramaria* sub. *Laeticolora*, *Ramaria* Sub. *Ramaria*, *Ramaria* Sub. *Echinormaria* and *Ramaria* sub. *Lentoramaria*. The species *Ramaria xizangensis* was grouped in *Ramaria* sub. *Laeticolora* along with *R. amyloidea* Marr & D.E. Stuntz, *R. celerivescens* Marr & D.E. Stuntz, and *R. claviramulata* Marr & D.E. Stuntz.

EU652352 Ramaria rubribrunnescens OSC 119676
KY354750 Ramaria rubribrunnescens OSC 66051
KY354749 Ramaria maculatipes OSC 112051
KT824242 Ramaria sp. KD-14-006
KY354743 Ramaria conjunctipes var. sparsiramosa OSC 119883
KC346861 Ramaria conjunctipes OSC 110613
EU846298 Ramaria araiospora OSC 108707
AJ408361 Ramaria subbotrytis MA-Fungi 48088
PP399157 Ramaria xizangensis CLZhao 31180
PP399158 Ramaria xizangensis CLZhao 31204
PP399156 Ramaria xizangensis CLZhao 31169
KX574472 Ramaria claviramulata WTU-F-043055
JX310392 Ramaria celerivirescens OSC 140471
EU837196 Ramaria amyloidea OSC 69891
OP177710 Ramaria patagonica 403
OP177713 Ramaria patagonica GM 19106
KJ655570 Gomphus ludovicianus TFB 14476
AJ292292 Gomphus clavatus MA-Fungi 48085
MK682678 Ramaria fennica AMB n. 17522
MN319564 Turbinellus cf floccosus MO 285170
MK493036 Ramaria flavescens AMB 17404
MG760613 Ramaria rasilisporoides MH-2013
KP658122 Ramaria stuntzii OSC 73315
EU669320 Ramaria coulterae p649i
MK169346 Ramaria rasilisporoides WTU-F-043029
MK493046 Ramaria pseudoflava AMB 17392
MK493035 Ramaria flava AMB 17393
KJ655554 Ramaria obtusissima TFB 14473
MK169351 Ramaria magnipes WTU-F-063057
JQ408240 Ramaria flavobrunnescens var. aromatica AGK 059
MT055910 Ramaria formosa AMB 18529
Ramaria sub. Laeticolora
JQ408221 Ramaria stricta var concolor AGK 011
MH856299 Ramaria stricta CBS:165.48
KP658148 Ramaria suecica OSC 115933
MK705858 Clavariadelphus sp HMAS 250466
JQ408236 Ramaria rubella f rubella AGK 049
MH860840 Ramaria apiculata var brunnea CBS:149.74
KP658128 Ramaria testaceoflava OSC 107885
JQ408230 Phaeoclavulina myceliosa AGK 035
MK796107 Phaeoclavulina flaccida AMB n. 17671
KY510818 Phaeoclavulina abietina u066
KJ140634 Kavinia alboviridis CFMR:DLL2011-131
KJ140598 Kavinia himantia CFMR:DLL2011-079
FJ596788 Lentaria byssiseda TENN61159
MF773634 Lentaria micheneri RRD6 TENN
MF992160 Ramaricium polyporoideum TENN:065654
EU669317 Ramaria rubella OSC 115946
KY626144 Ramaria botrytis f. musicolor ZT Myc 57160
KY626151 Ramaria botrytis AMB n. 18201
OP177707 Ramaria botrytis GM 19044
JX310410 Ramaria botrytis var. aurantiiramosa OSC 140667
KX574471 Ramaria botrytis var. aurantiiramosa WTU-F-043053
AF377059 Gautieria parksiana SNF236USA
MK169345 Ramaria fumosiavellanea WTU-F-063048
KY354738 Ramaria acrisiccescens OSC 112057
MK358820 Gloeocantharellus neoechinosporus GDGM75321
KP658144 Ramaria gelatiniaurantia OSC 65737
KX574480 Ramaria verlotensis WTU-F-063047
NR 155720 Ramaria luteovernalis MCVE 28637
AF442098 Ramaria aff. subtilis MA 48055
NR 137862 Ramaria admiratia TENN 69114
OP177715 Ramaria dendrophora GM 19094
OP177716 Ramaria dendrophora GM 20020
MN637783 Ramaria aurea AMB 18352
JQ408239 Ramaria foetida AGK 058
EU837197 Ramaria aurantiisiccescens OSC 104868
KP658130 Ramaria largentii OSC 67012
KP454028 Ramaria sandaracina var. sandaracina UBC F28386
OP177720 Ramaria flavinedulis GM 19035
OP177717 Ramaria flavinedulis GM 19056
OP177723 Ramaria inedulis 12648
OP177722 Ramaria inedulis GM 19047
MW683864 Lactarius sp PDD:113066
0.1

Figure 4. Phylogeny of species in the *Ramaria* generated by maximum likelihood based on ITS+LSU sequence data. Branches are labeled with maximum likelihood bootstrap ≥ 75% and Bayesian posterior probabilities ≥ 0.90, respectively. New species are in bold.

Taxonomy

Calocera ramaria **C.L. Zhao & H.M. Zhou, sp. nov.**
MycoBank No: 852565
Figs 5, 6

Holotype. CHINA, Xizang, Linzhi, Sejila Mountain National Forest Park, 29°64'N, 94°71'E, elev. 3852 m, gregarious on humus under *Abies*, 2 August 2023, CLZhao 31166 (SWFC).

Etymology. *Ramaria* (Lat.): refers to the ramal basidiomata of the specimens.

Diagnosis. Differed from other species in having ramal basidiomata, septate hyphae, usually 4-septate basidiospores (9.2–11 × 3.9–4.4 µm).

Fruiting body. Basidiomata stipitate, gregarious, bright orange when fresh, orange brown when dry, gelatinous when soaked, corneous when dry, ramal, repeatedly branched, apically blunt, up to 6.2 cm high; stipe 0.7–1 mm in diam, become orange to reddish brown corneous when dry.

Internal features. Marginal hyphae hyaline, smooth, thin-walled, septate, simple or branched, without clamp connections, 4–5.5 µm in diam; internal hyphae hyaline, smooth or scabrous, thin- to slightly thick-walled, interwoven, with nodose-septa, without clamp connections, 2–3 µm in diam; hyphidia hyaline, smooth, thin-walled, with a simple septum at base, occasionally terminally branched; basidia hyaline, thin-walled, subclavate to clavate, without basal clamp connection, 23–31 × 2–4 µm; basidiospores hyaline, smooth, thin-walled, oblong-ellipsoid to navicular, straight or curved, apiculate, usually 4-septate when mature, occasionally 5-septate, (9.1–)9.2–11(–11.6) × (3.5–)3.9–4.4(–4.7) µm, L = 10.18 µm, W = 4.19 µm, Q = 2.43 (n = 30/1).

Ceraceomyces rhizomorphus **C.L. Zhao & H.M. Zhou, sp. nov.**
MycoBank No: 852584
Figs 7, 8

Holotype. CHINA, Xizang, Linzhi, Sejilashan National Forest Park, 29°64'N, 94°71'E, elev. 3848 m, on the fallen branch of *Abies*, 2 August 2023, CLZhao 31188 (SWFC).

Etymology. *Rhizomorphus* (Lat.): refers to the basidiomata with rhizomorphs.

Diagnosis. Differed from other species in having merulioid, cream to yellowish basidiomata, generative hyphae with clamp connections, cylindrical basidiospores (4.7–6.2 × 1.8–2.3 µm).

Fruiting body. Basidiomata resupinate, adnate, smooth to tuberculate when fresh, merulioid upon drying, without odor or taste when fresh, up to 6 cm long, 2 cm wide, 100–200 µm thick. Hymenial surface merulioid, cream to yellowish when fresh, turn to orange yellow upon drying. Margin sterile, white, with rhizomorphs.

Hyphal structure. Hyphal system monomitic, generative hyphae with clamp connections, colorless, thin- to slightly thick-walled, branched, interwoven, 3.5–7 µm in diameter, IKI–, CB–; tissues turn black in KOH.

Hymenium. Cystidia and cystidioles absent; basidia narrowly clavate to clavate, in a dense palisade, with 4 sterigmata and a basal clamp connection,



Figure 5. Basidiomata and microscopic structures of *Calocera ramaria* (holotype, CLZhao 31166, holotype) **A, B** basidiomata **C** a section of hymenium **D** basidiospores **E** marginal hyphae **F** internal hyphae. Scale bars: 1 cm (**A, B**); 10 μm (**C–F**).

16–19 × 3.5–4 μm; basidioles dominant, similar to basidia in shape, but slightly smaller.

Spores. Basidiospores cylindrical, with suprahilar depression, colorless, smooth, thin-walled, IKI–, CB–, (4.2–)4.7–6.2(–6.4) × (1.5–)1.8–2.3(–2.4) μm, L = 5.49 μm, W = 2.05 μm, Q = 2.66–2.68 (n = 60/2).



Figure 6. Microscopic structures of *Calocera ramaria* (holotype, CLZhao 31166) **a** basidiospores **b** basidia with basidiospores. Scale bars: 5 μm (**a**); 10 μm (**b**).

Additional specimens examined (*paratypes*). China. Xizang, Linzhi, Sejila Mountain National Forest Park, 29°64'N, 94°71'E, elev. 3848 m, on the trunk of *Abies*, 2 August 2023, CLZhao 31153 (SWFC); CLZhao 31154 (SWFC); CLZhao 31161 (SWFC); CLZhao 31202 (SWFC); on the fallen branch of *Abies*, 2 August 2023, CLZhao 31184 (SWFC); CLZhao 31185 (SWFC); CLZhao 31197 (SWFC).



Figure 7. Basidiomata of *Ceraceomyces rhizomorphus* **A, C** CLZhao 31188 (holotype) **B, D** CLZhao 31185.



Figure 8. Microscopic structures of *Ceraceomyces rhizomorphus* (holotype, CLZhao 31216) **a** basidiospores **b** basidia **c** basidioles **d** a section of hymenium. Scale bars: 5 μm (**a**); 10 μm (**b–d**).

Leptosporomyces linzhiensis **C.L. Zhao & H.M. Zhou, sp. nov.**

MycoBank No: 852585

Figs 9, 10

Holotype. CHINA, Xizang, Linzhi, Sjilashan Forest Park, 29°64'N, 94°71'E, elev. 3848 m, on fallen trunk of *Abies*, 2 August 2023, CLZhao 31183 (SWFC).

Etymology. *Linzhiensis* (Lat.): refers to the locality (Xizang) of the type specimens.

Diagnosis. Differed from other species in having white basidiomata, monomitic hyphal system, cylindrical to oblong ellipsoid basidiospores (3.8–4. × 1.7–2 µm).

Fruiting body. Basidiomata resupinate, athelioid, membranous upon drying, without odor or taste when fresh, up to 10 cm long, 4 cm wide, 200 µm thick. Hymenial surface smooth to cracked, white with pink tint when fresh, turning to yellowish cream upon drying. Margin sterile, white, fimbriate.

Hyphal structure. Hyphal system monomitic, generative hyphae with clamp connections, colorless, thin- to slightly thick-walled, branched, interwoven, 2–5 µm in diameter, IKI–, CB–; tissues turn black in KOH.

Hymenium. Hyphal system monomitic, generative hyphae with clamp connections, colorless, thin-walled, branched, interwoven, 2–3.5 µm in diameter, IKI–, CB–. Basidia clavate, with 4 sterigmata and a basal clamp connection, 11.5–13.5 × 3.2–3.8 µm.

Spores. Basidiospores cylindrical to oblong ellipsoid, colorless, smooth, thin-walled, IKI–, CB–, (3.5–)3.8–4.3(–4.7) × (1.7–)1.7–2(–2.3) µm, L = 4.02 µm, W = 1.88 µm, Q = 1.95–2.18 (n = 90/3).

Additional specimens examined (*paratypes*). CHINA, Xizang, Linzhi, Sjilashan Forest Park, 22°57'N, 103°42'E, elev. 2100 m, on fallen trunk of *Abies*, 2 August 2023, CLZhao 31174 (SWFC); on fallen trunk of *Abies*, 2 August 2023, CLZhao 31187 (SWFC); on fallen trunk of *Abies*, 2 August 2023, CLZhao 31190 (SWFC).



Figure 9. Basidiomata of *Leptosporomyces linzhiensis* (holotype, CLZhao 31183). Scale bars: 1 cm (**A**); 1 mm (**B**).



Figure 10. Microscopic structures of *Leptosporomyces linzhiensis* (holotype, CLZhao 31183) **a** basidiospores **b** basidia **c** basidioles **d** A section of hymenium. Scale bars: 5 μm (**a**); 10 μm (**b–d**).

Ramaria xizangensis **C.L. Zhao & H.M. Zhou, sp. nov.**

MycoBank No: 852586

Figs 11, 12

Holotype. China, Xizang, Linzhi, Sejila Mountain National Forest Park, 29°64'N, 94°71'E, elev. 3850 m, gregarious on the humus under *Abies*, 2 August 2023, CLZhao 31169 (SWFC).

Etymology. *Xizangensis* (Lat.): refers to the locality (Xizang) of the type specimens.

Diagnosis. Differed from other species in having flesh pink basidiomata, monomitic hyphal system, generative hyphae with clamp connections, ellipsoid to cylindrical, densely warted basidiospores (9.7–11.8 × 3.9–4.9 µm).

Fruiting body. Basidiomata solitary to gregarious, with 8 cm high × 6 cm wide at the widest point, repeat branched dichotomously in 4–5 ranks, flesh pink when fresh, become clay buff with dry; apices obtuse, orange yellow when fresh, becoming fuscous when dry. Stipe ≥ 3 cm high, compound to fasciculate in groups of 5, emerging from a common base, concolorous with the branches.

Hyphal structure. Hyphal system monomitic, generative hyphae with clamp connections, branched, walls smooth and hyaline; basal stem with tramal hyphae 4–7 µm wide and inflated ones up to 10 µm, occasionally branched, thin-walled, parallel arranged, hyaline; tramal hyphae of branches 3–4 µm wide.

Hymenium. Hymenium all along the basidiomata. Basidia clavate, in a dense palisade, with 4 sterigmata and a basal clamp connection. Basidioles elongated clavate, smooth, hyaline, contents homogeneous, 23.5–34 × 6–7 µm.

Spores. Basidiospores ellipsoid to cylindrical, densely warted, with 1–2 several guttulae, IKI–, CB–, 9.7–11.8(–12.5) × (3.8–)3.9–4.9(–5.1) µm, L = 10.69 µm, W = 4.29 µm, Q = 2.49 (n = 30/1).

Additional specimens examined (*paratypes*). CHINA, Xizang, Linzhi, Sejila Mountain National Forest Park, 29°67'N, 94°74'E, elev. 3850 m, gregarious on the humus under *Abies*, 2 August 2023, CLZhao 31180 (SWFC); on ground in forest of *Abies*, 2 August 2023, CLZhao 31204 (SWFC).



Figure 11. Basidiomata of *Ramaria xizangensis* (holotype, CLZhao 31169). Scale bars: 1 cm (**A, B**).



Figure 12. Microscopic structures of *Ramaria xizangensis* (CLZhao 31169, holotype) **a** basidiospores **b** basidia **c** a section of hymenium. Scale bars: 5 µm (**a**); 10 µm (**b, c**).

Discussion

Wood decay fungi encompasses the vast group of aphyllophoroid fungi with corticioid, prioid or jelly form of basidiomata (Herter 1910). This classification has historically been used to define the different families of Basidiomycetes. However, molecular studies have revealed that many of these fungi are distributed across various orders within the Basidiomycetes, including the likes

of Amylocorticiales, Atheliales, Dacrymycetales, and Gomphales (Kirk et al. 2018; Wei et al. 2022). As a result, further research is needed to elucidate the relationships and morphological variability of these taxa through phylogenetic analysis.

The Xizang Autonomous Region, situated in the southwest of China, is renowned as one of the most bio-diverse regions in the country. This is attributed to its complex topography and diverse ecosystems, making it a focal point for fungal biodiversity in China. Recently, studies focusing on fungal diversity and the ecology of Basidiomycota in Xizang were carried out (Ke 2016; Pubu et al. 2016; Wang et al. 2023). According to the study (Pubu et al. 2016), 1733 species were collected in Xizang. The fungal research indicated that Sejila Mountain National Forest Park is predominantly composed of spruce and fir trees, which provide an ideal habitat for a rich diversity of macrofungi species to flourish (Zhao and Li 1987). In our study, four species were found from Xizang, *Calocera ramaria*, *Ceraceomyces rhizomorphus*, *Leptosporomyces linzhiensis*, and *Ramaria xizangensis*.

Calocera is characterized by its yellow, gelatinous basidiomata, resembling *Dacrymyces*. However, *Dacrymyces* displays a broader range of basidiomata forms, including pulvinate, discoid, turbinate, spathulate, flabellate, and cylindrical shapes (Shirouzu et al. 2009; Fan et al. 2021), whereas *Calocera* exhibits branched, dendroid basidiomata. Our results have further confirmed that our newly discovered species features ramal basidiomata and clusters phylogenetically with *Calocera* species, placing it within the genus *Calocera*. In Xizang, two species have been identified, *C. ramaria* and *C. tibetica*, but the latter has wider basidiospores (5–6 µm vs. 3.9–4.4 µm, Fan et al. 2021). In our phylogenies, *C. viscosa* and *C. cornea* were related to *C. ramaria* (Fig. 1); however, *C. viscosa* has 1-septate mature basidiospores, and *C. cornea* differs from *C. ramaria* by its distinctly larger basidiospores (7–10 × 3–4.5 µm vs. 9.2–11 × 3.9–4.4 µm) with one septum (McNabb 1965; Shirouzu et al. 2009).

Previous research has highlighted the polyphyly of *Ceraceomyces* (Chikowski 2016; Yuan et al. 2023), and seven species are retained in *Ceraceomyces*. However, it is worth noting that authentic specimens and DNA data are lacking for *Ceraceomyces* species. Phylogenetically, *C. rhizomorphus* formed a sister group with *C. yunnanensis* and *C. borealis*, but *C. yunnanensis* has smaller basidiospores (3–4 × 1–1.5 µm vs. 4.7–6.2 × 1.8–2.3 µm, Yuan et al. 2023) and *C. borealis* has larger basidiospores (6–8 × 1.8–2 µm vs. 4.7–6.2 × 1.8–2.3 µm, Bernicchia and Gorjón 2010).

Ceraceomyces rhizomorphus and *C. tessulatus* had similar yellowish basidiomata with rhizomorphs when fresh, while *C. tessulatus* has ellipsoid and larger basidiospores (6–8 × 3.5–4.5 µm vs. 4.7–6.2 × 1.8–2.3 µm, Bernicchia and Gorjón 2010). Three known species, *C. bizonatus*, *C. reidii*, and *C. simulans* also distributed in Asia. However, *C. bizonatus* has shorter basidiospores (2.5–3.3 µm vs. 4.7–6.2 µm, Bernicchia and Gorjón 2010); *C. reidii* has larger basidiospores (11.5–15 × 4.5–6 µm vs. 4.7–6.2 × 1.8–2.3 µm, Bernicchia and Gorjón 2010); *C. simulans* has longer basidiospores (6–7 µm vs. 4.7–6.2 µm, Bernicchia and Gorjón 2010).

Leptosporomyces linzhiensis is similar to *L. thindii* in having white basidiomata and being distributed in Asia, but the latter has wider basidiospores (Prasher 2015). *Leptosporomyces linzhiensis* sisters to *L. septentrionalis* by its white ba-

sidiomata, and cylindrical basidiospores, but the latter has slightly shorter basidiospores (3–4 µm vs. 3.8–4.3 µm), and 2–4 basidia (Prasher 2015). *Leptosporomyces linzhiensis* is easily confused with *L. roseus* in, but the latter has shorter basidiospores (2–2.5 µm vs. 3.8–4.3 µm, Prasher 2015). *Leptosporomyces fuscostratus* has a broad distributional range in the northern hemisphere, but it has wider basidiospores (2–2.8 µm vs. 1.7–2 µm, Yurchenko and Wołkowycki 2022).

In our phylogeny, *Ramaria* is paraphyletic, which included four clades, *R.* sub. *Laeticolora* and *R.* sub. *Lentoramaria*, *R.* sub. *Ramaria* and R. sub. *Echinormaria*. *Ramaria xizangensis* was clustered in *Ramaria* sub. *Laeticolora* with *Ramaria amyloidea*, *R. celerivirescens* and *R. claviramulata*. However, *R. celerivirescens* has slightly wider basidiospores (4–6 µm vs. 3.9–4.9 µm, Marr and Stuntz 1973). *R. claviramulata* has cream to brownish white basidiomata. *Ramaria xizangensis* is similar to *R. indoyunnaniana* in having pink basidiomata and being distributed in Yunnan, but the latter has shorter basidiospores (7.2–8.3 µm vs. 9.7–11.8 µm, Petersen and Zang 1986).

According to our field inventory, the four Chinese new species were found in alpine zone near the Sejila Mountain, and the coniferous forest dominant by *Abies* at high altitude with cold and humid environments. Previously, numerous new species have been found in Southwest China (Dai 2022; Zhao et al. 2023), and the present paper confirms the fungal diversity is very rich in the montane forests of Southeast Xizang.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The research was supported by the National Natural Science Foundation of China (Project No. 32170004); the Highlevel Talents Program of Yunnan Province (YNQR-QNRC-2018-111); the Scientific Research Fund of Yunnan Provincial Department of Education (2024J0668); Forestry Innovation Programs of Southwest Forestry University (Grant No: LXXK-2023M07).

Author contributions

Data curation: ZHM, ZCL,WF. Formal analysis: ZXC. Methodology: ZHM, ZXC, LJT. Software: ZXC, LJT. Writing - original draft: ZHM, ZCL. Writing - review and editing: ZCL, WF.

Author ORCIDs

Hong-Min Zhou https://orcid.org/0000-0002-0724-5815
Xun-Chi Zhang https://orcid.org/0000-0003-3887-0979
Fang Wu https://orcid.org/0000-0002-1455-6486
Chang-Lin Zhao https://orcid.org/0000-0002-8668-1075

Data availability

All of the data that support the findings of this study are available in the main text.

References

Anonymous (1969) Flora of British Fungi. Colour Identification Chart. Her Majesty's Stationery Office, London.

Bao Y (2020) Mycoflora Biodiversity and Genetic Variation of Macrofungi in Alpine Meadow of Tibet. Northeast Normal University.

Bernicchia A, Gorjón SP (2010) Fungi Europaei 12. Corticiaceae s.l. Edizioni Candusso, Alassio, Italy, 1008 pp.

Binder M, Larsson KH, Matheny PB, Hibbett DS (2010) Amylocorticiales ord. nov. and Jaapiales ord. nov.: Early diverging clades of Agaricomycetidae dominated by corticioid forms. Mycologia 102(4): 865–880. https://doi.org/10.3852/09-288

Chikowski R, Larsson KH, Gibertoni TB (2016) *Ceraceomyces atlanticus* (Amylocorticiales, Basidiomycota), a new species from the Atlantic Rain Forest, Brazil. Phytotaxa 296(1): 73–80. https://doi.org/10.11646/phytotaxa.296.1.5

Coker WC (1923) The *Clavarias* of the United States and Canada. Chapel Hill, 209 pp.

Corner EJH (1950) A monograph of *Clavaria* and allied genera. Annals of Botany Memoirs 1: 1–740.

Cui BK, Li HJ, Ji X, Zhou JL, Song J, Si J, Yang ZL, Dai YC (2019) Species diversity, taxonomy and phylogeny of Polyporaceae (Basidiomycota) in China. Fungal Diversity 97(1): 137–392. https://doi.org/10.1007/s13225-019-00427-4

Doty MS (1944) *Clavaria*, the species known from Oregon and the Pacific Northwest.

El-Gharabawy HM, Leal-Dutra CA, Griffith GW (2021) *Crystallicutis* gen. nov. (Irpicaceae, Basidiomycota), including *C. damiettensis* sp. nov., found on *Phoenix dactylifera* (date palm) trunks in the Nile Delta of Egypt. Fungal Biology 125(6): 447–458. https://doi.org/10.1016/j.funbio.2021.01.004

Exeter RL, Norvell L, Cazares E (2006) *Ramaria* of the Pacific Northwestern United States. United States Department of Interior, Bureau of Land Management. Salem, Oregon USA, 156 pp.

Fan LF, Wu YD, Wu F, Dai YC (2021) *Calocera tibetica* sp. nov. (Dacrymycetaceae, Dacrymycetales) from southwestern China. Phytotaxa 500: 133–141. https://doi.org/10.11646/phytotaxa.500.2.6

Fisher MC (1931) A comparative morphological study of certain species of the Dacryomycetaceae. The Proceedings 38(1): 115–125.

Gäumann EA (1953) The Fungi: a description of their morphological features and evolutionary development, (Translated by Wynd FL). Hafner Publishing, New York and London, 420 pp.

Giachini AJ, Hosaka K, Nouhra E, Spatafora J, Trappe JM (2010) Phylogenetic relationships of the Gomphales based on nuc-25S-rDNA, mit-12S-rDNA, and mit-atp6-DNA combined sequences. Fungal Biology 114(2–3): 224–234. https://doi.org/10.1016/j.funbio.2010.01.002

Hall TA (1999) Bioedit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.

Hanif M, Khalid AN, Exeter RL (2019) *Ramaria flavescentoides* sp. nov. with clamped basidia from Pakistan. Mycotaxon 134: 399–406. https://doi.org/10.5248/134.399

Herter WGF (1910) Autobasidiomycetes. Kryptogamen-Flora der Mark Brandenburg 6: 1–192.

Hodkinson BP, Moncada B, Lücking R (2014) Lepidostromatales, a new order of lichenized fungi (Basidiomycota, Agaricomycetes), with two new genera, *Ertzia* and *Sul-*

zbacheromyces, and one new species, *Lepidostroma winklerianum*. Fungal Diversity 64(1): 165–179. https://doi.org/10.1007/s13225-013-0267-0

Hosaka K, Bates ST, Beever RE, Castellano MA, Colgan W III, Dominguez LS, Trappe JM (2006) Molecular phylogenetics of the gomphoidphalloid fungi with an establishment of the new subclass Phallomycetidae and two new orders. Mycologia 98(6): 949–959. https://doi.org/10.1080/15572536.2006.11832624

Humpert AJ, Muench EL, Giachini AJ, Castellano MA, Spatafora JW (2001) Molecular phylogenetics of *Ramaria* and related genera: evidence from nuclear large subunit and mitochondrial small subunit rDNA sequences. Mycologia 93: 465–477. https://doi.org/10.2307/3761733

Jülich W (1972) Monographie der Athelieae (Corticiaceae, Basidiomycetes). Willdenowia Beiheft 7: 1–283.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14(6): 587–589. https://doi.org/10.1038/nmeth.4285

Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. Briefings in Bioinformatics 20(4): 1160–1166. https://doi.org/10.1093/bib/bbx108

Ke ZD (2016) Studies on macrofungi diversity in Shergyla Mountain National Nature Reserve in Tibet. Jilin Agricultural University.

Kirk PM, Ainsworth GC, Cannon PF, Minter DW (2008) Ainsworth and Bisby's dictionary of fungi, 10th edn. CAB International, UK. https://doi.org/10.1079/9780851998268.0000

Knudson AG (2012) The genus *Ramaria* in Minnesota. Master of Science thesis, University of Minnesota.

Larsson KH (2007) Re-thinking the classification of corticioid fungi. Mycological Research 111(9): 1040–1063. https://doi.org/10.1016/j.mycres.2007.08.001

Larsson KH, Larsson E, Kõljalg U (2004) High phylogenetic diversity among corticoid homobasidiomycetes. Mycological Research 108(9): 983–1002. https://doi.org/10.1017/S0953756204000851

Liu S, Chen YY, Sun YF, He XL, Song CG, Si J, Liu DM, Gates G, Cui BK (2023) Systematic classification and phylogenetic relationships of the brown-rot fungi within the Polyporales. Fungal Diversity 118(1): 1–94. https://doi.org/10.1007/s13225-022-00511-2

Lowy B (1971) Tremellales. Flora Neotropica 6(8): 1–153.

Marr CD, Stuntz DE (1973) *Ramaria* of Western Washington. Bibliotheca Mycologica, Germany, 38 pp.

Martin GW (1948) New of noteworthy tropical fungi. IV. Lloydia 11(2): 111–122.

McNabb RFR (1965) Taxonomic studies in the Dacrymycetaceae 2. *Calocera* (Fries) Fries. New Zealand Journal of Botany 3: 31–58. https://doi.org/10.1080/002882 5X.1965.10428712

Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In Proceedings of the Gateway Computing Environments Workshop, New Orleans, 8 pp. https://doi.org/10.1109/GCE.2010.5676129

Nees Esenbeck, CDG (1817) System der Pilze und Schwämme, 334 pp.

Peng YB, Liu B, Li F (1992) Flora Fungorum Sinicorum Vol. 2. Tremellales et Dacrymycetales. Science Press, Beijing, 151 pp.

Petersen JH (1996) Farvekort. The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve, 6 pp.

Petersen RH (1981) Ramaria subgenus Echinoramaria. Bibliotheca Mycologia 261 pp.

Petersen RH, Zang M (1986) New or interesting clavarioid fungi from Yunnan, China. Plant Diversity 8(3): 281–294.

Prasher IB (2015) Wood-rotting non-gilled Agaricomycetes of Himalayas. Fungal Diversity Research Series, Chiang Rai, 653 pp. https://doi.org/10.1007/978-94-017-9858-7

Pubu CR, Wang M, Liu XY (2016) A review of investigation of fungi in Tibet. Mycosystema 35(9): 1025–1047. https://doi.org/10.13346/j.mycosystema.150158

Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19(12): 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Ryvarden L, Gilbertson RL (1993) European polypores. Part 1. Synopsis Fungorum 6: 1–387.

Shirouzu T, Hirose D, Tokumasu S (2007) Sequence analyses of the 28S rRNA gene D1/D2 region suggest *Dacrymyces* (Heterobasidiomycetes, Dacrymycetales) is polyphyletic. Mycoscience 48(6): 388–394. https://doi.org/10.1007/S10267-007-0378-0

Shirouzu T, Hirose D, Tokumasu S (2009) Taxonomic study of the Japanese Dacrymycetes. Persoonia 23(1): 16–34. https://doi.org/10.3767/003158509X468443

Shirouzu T, Hosaka K, Nam KO, Weir BS, Johnston PR, Hosoya T (2017) Phylogenetic relationships of eight new Dacrymycetes collected from New Zealand. Persoonia 38(1): 156–169. https://doi.org/10.3767/003158517X695280

Sulistyo BP, Larsson KH, Haelewaters D, Ryberg M (2021) Multigene phylogeny and taxonomic revision of Atheliales s.l.: Reinstatement of three families and one new family, Lobuliciaceae fam. nov. Fungal Biology 125(3): 239–255. https://doi.org/10.1016/j.funbio.2020.11.007

Wang SH, Li GJ, Phurbu D, He MQ, Zhang MZ, Zhu XY, Li JX, Zhao RL, Cao B (2023) Four new species of *Russula* from the Xizang Autonomous Region and other provinces of China. Mycology 15(2): 1–28. https://doi.org/10.1080/21501203.2023.2265667

Wei CL, Chen CC, He SH, Wu SH (2022) *Dendrocorticiopsis orientalis* gen. et sp. nov. of the Punctulariaceae (Corticiales, Basidiomycota) revealed by molecular data. MycoKeys 90: 19–30. https://doi.org/10.3897/mycokeys.90.84562

White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: MA, Gefand DH, Sninsky JJ, White MJT (Eds) PCR Protocols: a guide to methods and applications; Academic Press, San Diego, 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, et al. (2020) Outline of fungi and fungus-like taxa. Mycosphere 11(1): 1160–1456. https://doi.org/10.5943/mycosphere/11/1/8

Wu SH, Shih K, Yu SY (2011) *Calocera bambusicola* sp. nov. and *C. sinensis* newly recorded from Taiwan. Mycotaxon 115(1): 163–169. https://doi.org/10.5248/115.163

Yang Y, Jiang Q, Li Q, Yang J, Cha L, Cheng L, Yang S, Zhao C, Zhou H (2023) Molecular systematics and taxonomic analyses of three new wood-inhabiting fungi of *Hyphoderma* (Hyphodermataceae, Basidiomycota). Journal of Fungi 9(11): 1044. https://doi.org/10.3390/jof9111044

Yuan Q, Luo KY, Zhang Y, Zhao CL (2023) Morphological characteristics and phylogenetic analyses revealed three new wood-inhabiting fungi (Agaricomycetes, Basidiomycota) in Southern China. Phytotaxa 592(3): 179–195. https://doi.org/10.11646/phytotaxa.592.3.1

Yurchenko K, Wołkowycki M (2022) New species of corticioid fungi (Basidiomycota) for Poland found in Białowieża Primeval Forest in 2018–2020. Acta Mycologica 58: 1–18. https://doi.org/10.5586/am.577

Zamora JC, Ekman S (2020) Phylogeny and character evolution in the Dacrymycetes, and systematics of Unilacrymaceae and Dacryonaemataceae fam. nov. Persoonia 44(1): 161–205. https://doi.org/10.3767/persoonia.2020.44.07

Zhao LF, Li XG (1987) Partial fungal list of Nyingchi region, Milin, Tibet. Journal of Southwest Forestry College 7(1): 55–58. [Natural Science]

Zhao CL, Qu MH, Huang RX, Karunarathna SC (2023) Multi-gene phylogeny and taxonomy of the wood-rotting fungal genus *Phlebia* sensu lato (Polyporales, Basidiomycota). Journal of Fungi 320(9): 1–41. https://doi.org/10.3390/jof9030320